Mach, K., (367) 69
Makosza, M., (367) 95
Martinet, P., (367) 1
Masamune, S., (367) 39
Masters, A.F., (367) 371
Mathur, P., (367) 243
Matt, D., (367) 117
Mavunkal, I.J., (367) 243
Mita, S., (367) 81
Mori, N., (367) 81
Müller, J., (367) 133
Mukkanti, K., (367) C29

Neumann, F., (367) 175
Nieger, M., (367) 47
Normant, J.-R., (367) 1
Nuber, B., (367) 197

Ochs, M., (367) 343
Oder, K., (367) 249
Okuda, J., (367) C1
Ono, I., (367) 81
Orrell, K.G., (367) 275
Orsini, F., (367) 375
Osborne, A.G., (367) 275
Ostrowski, S., (367) 95

Park, J., (367) 39
Pelizzoni, F., (367) 375
Périchon, J., (367) 347
Pörschke, K.R., (367) 233
Popod'ko, N.R., (367) 205
Pozdeeva, A.A., (367) 205

Reisner, G.M., (367) 77
Ricalton, A., (367) C21
Ridder, D.J.A. de, (367) 55
Rieker, J., (367) 107
Roder, K., (367) 321, 339
Rogers, G.W., (367) 101
Rollin, Y., (367) 347
Rubinsztajn, S., (367) 27
Rugmini, V., (367) 243

Sauvêtre, R., (367) 1
Sawitzki, G., (367) 19
Schaaf, P.A. van der, (367) 55
Schulte, H.-G., (367) 359
Scordia, H., (367) 143
Seevogel, K., (367) 233
Sellmann, D., (367) 161
Shaver, A., (367) C5
Šik, V., (367) 275
Skelton, B.W., (367) C25
Sonn, I., (367) 133
Subba Rao, Y.V., (367) C29
Süss-Fink, G., (367) 175

Teuben, J.H., (367) C18
Theiler, H.G., (367) 291
Tolstikov, G.A., (367) 205

Vallarino, L.M., (367) 375
Van Koten, G., (367) 55
Van der Schaaf, P.A., (367) 55
Varga, V., (367) 69
Vogler, C., (367) 107
Vondrák, T., (367) 69

Wachter, J., (367) 197
Watson, A.A., (367) C25
Weike, Z., (367) C9
Werner, H., (367) 321, 339
White, A.H., (367) C25
Whiteley, M.W., (367) C21
Wilbur, D.S., (367) 259

Zaworotko, M.J., (367) 267
Zhdanov, S.I., (367) 205
Ziegler, M.L., (367) 197
Zwettler, R., (367) C15

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 367 (1989)

SUBJECT INDEX

Aluminium

$(C_5H_5)_2ZrCl(AlCl_4)$, crystal and molecular structure and unusual hydrocarbon solubility properties of; relevance to Ziegler–Natta polymerization chemistry (M.V. Gaudet, M.J. Zaworotko, T.S. Cameron, A. Linden), (367) 267

Boron

Boron hydride insertion into Zr–C bond of dimeric (η^2-formaldehyde)zirconocene (G. Erker, U. Hoffmann, R. Zwettler, G. Krüger), (367) C15

(1,1-Dichloroalkyl)lithium, reactions with boronic acid of; facile synthesis of tertiary carbinols (B. Junchai, Z. Weike, D. Hongxun), (367) C9

2,4,6,8,9,10-Hexachloro-2,4,6,8,9,10-hexaboraadamantane (W. Haubold, W. Keller, G. Sawitzki), (367) 19

Methyl 4-tri-n-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Chromium

η^6-Arenetricarbonylchromium(0) π-complexes, failing nucleophilic substitution of hydrogen in (S. Ostrowski, M. Mąkosza), (367) 95

Benchrotrenic alcohols, asymmetric synthesis with baker's yeast of (J. Gillois, G. Jaouen, D. Buisson, R. Azerad), (367) 85

Group 6 metal tetracarbonyl complexes, *cis*-$[M(CO)_4\{(C_5H_4ECH_3)_2Fe\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1′-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Tricarbonyl(boat *p*-xylene)chromium complexes, ab initio calculations of the charge densities in (I. Ono, S. Mita, S. Kondo, N. Mori), (367) 81

Cobalt

$(\eta^5\text{-}C_5Me_5)_2CoRh(\mu\text{-}CO)_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197

Dicarbonylcyclopentadienylmethylthiolatoiron(II), reactions with monocyanoacetylene of; crystal structures of $[FeC(O)C(CN)=CH(SCH_3)(\eta^5\text{-}C_5H_5)(CO)]$ and of $[\mu\text{-}\eta^2\text{-}Fe(C=CCN)(\eta^5\text{-}C_5H_5)(CO)_2][Co(CO)_3]_2$ (R. Kergoat, M.M. Kubicki, L.C. Gomes de Lima, H. Scordia, J.E. Guerchais, P. L'Haridon), (367) 143

Tetrakis- and hexakis-(trimethylsilyl)cobaltocenium ions, ring rotation hindrance in (J. Okuda), (367) C1

Trimethylphosphinecobalt(I) complexes: reaction with cyclopentatriene and cyclooctatetraene; crystal structure of $[Co(\eta^4\text{-}C_7H_8)(PMe_3)_3]BPh_4$ (L.C. Ananias de Carvalho, M. Dartiguenave, Y. Dartiguenave), (367) 187

Copper

Cu^I-macrocyclic tridentate amine systems, reactions with CO, C_2H_4 and cyclic olefins of (P. Chaudhuri, K. Oder), (367) 249

Germanium

Tetrakis(2,6-diisopropylphenyl)digermene, synthesis of and its reductive cleavage to the corresponding digermenyllithium(DME) complex (J. Park, S.A. Batcheller, S. Masamune), (367) 39

Iron

1,6-Bis(triphenylmethyl)biferrocenium triiodide, electron transfer in (T.-Y. Dong, T.-L. Hsu), (367) 313

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(CO)_6Fe_2(\mu_3\text{-}E)_2M(PPh_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

$[(C_5H_4Me)FeCO]_3NO$ and $[(C_5Me_5)FeCO]_3NO$, synthesis and structure of (J. Müller, I. Sonn, T. Akhnoukh), (367) 133

Dicarbonylcyclopentadienylmethylthiolatoiron(II), reactions with monocyanoacetylene of; crystal structures of $[\overline{FeC(O)C(CN)=CH(SCH_3)}(\eta^5\text{-}C_5H_5)(CO)]$ and of $[\mu\text{-}\eta^2\text{-}Fe(C{=}CCN)(\eta^5\text{-}C_5H_5)(CO)_2][Co(CO)_3]_2$ (R. Kergoat, M.M. Kubicki, L.C. Gomes de Lima, H. Scordia, J.E. Guerchais, P. L'Haridon), (367) 143

β-Keto phosphines derived from ferrocene; syntheses and structures of $[Ph_2PCH_2C(O)(\eta^5\text{-}C_5H_4)Fe(\eta^5\text{-}C_5H_5)](L^1)$ and *trans*-$[PdCl_2L^1_2]$ (P. Braunstein, T.M. Gomes Carneiro, D. Matt, F. Balegroune, D. Grandjean), (367) 117

Ruthenium dinuclear cluster with N- and O-containing bridges, reactions with CO_2 of (F. Neumann, G. Süss-Fink), (367) 175

Lithium

(1,1-Dichloroalkyl)lithium, reactions with boronic acid of; facile synthesis of tertiary carbinols (B. Junchai, Z. Weike, D. Hongxun), (367) C9

$Et_3SiCF{=}CFM$ (M = Li, ZnBr), preparation and reactivity of (P. Martinet, R. Sauvêtre, J.-F. Normant), (367) 1

Tetrakis(2,6-diisopropylphenyl)digermene, synthesis of and its reductive cleavage to the corresponding digermenyllithium(DME) complex (J. Park, S.A. Batcheller, S. Masamune), (367) 39

Magnesium

Grignard reactions of 1-phenylethyl chloride with aldehydes and ketones, reactivity of benzyl Grignard reagent (C. Bernardon), (367) 11

Mercury

Methyl 4-tri-n-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Metallocenes

1,6-Bis(triphenylmethyl)biferrocenium triiodide, electron transfer in (T.-Y. Dong, T.-L. Hsu), (367) 313

Boron hydride insertion into Zr–C bond of dimeric (η^2-formaldehyde)zirconocene (G. Erker, U. Hoffmann, R. Zwettler, G. Krüger), (367) C15

Dicyclopentadienyltitanium dihalides, electronic structure as indicated by UV photoelectron spectrocopy of (T. Vondrák, K. Mach, V. Varga), (367) 69

Dicyclopentadienyltricarbonyldirhodium, search for ^{13}C equilibrium isotope effects in (S. Berger, M. Ochs), (367) 343

Group 6 metal tetracarbonyl complexes, *cis*-$[M(CO)_4\{(C_5H_4ECH_3)_2Fe\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1′-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

β-Keto phosphines derived from ferrocene; syntheses and structures of $[Ph_2PCH_2C(O)(\eta^5\text{-}C_5H_4)Fe(\eta^5\text{-}C_5H_5)](L^1)$ and *trans*-$[PdCl_2L^1_2]$ (P. Braunstein, T.M. Gomes Carneiro, D. Matt, F. Balegroune, D. Grandjean), (367) 117

Palladium(0)-mediated reactions of $BrZnCH(CH_3)COO^tBu$ with vinyl and aryl triflates, influence of catalyst on (F. Orsini, F. Pelizzoni, L.M. Vallarino), (367) 375

Tetrakis- and hexakis-(trimethylsilyl)cobaltocenium ions, ring rotation hindrance in (J. Okuda), (367) C1

Molybdenum

Group 6 metal tetracarbonyl complexes, *cis*-[M(CO)$_4${(C$_5$H$_4$ECH$_3$)$_2$Fe}] (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1′-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Ligand-centred coupling of alkynyl radical cations [Mo(C≡CR)(Ph$_2$PCH$_2$CH$_2$PPh$_2$)(η-C$_7$H$_7$)]$^+$ (R = Ph or Bu^n): crystal structure of [Mo$_2$(Ph$_2$PCH$_2$CH$_2$PPh$_2$)$_2$(η-C$_7$H$_7$)$_2$(μ-C$_4$R$_2$)][PF$_6$]$_2$ (R = Ph) (R.L. Beddoes, C. Bitcon, A. Ricalton, M.W. Whiteley), (367) C21

Molybdenum(II) and tungsten(II) cationic seven-coordinate complexes, synthesis of (P.K. Baker, M. Bamber, G.W. Rogers), (367) 101

Nickel

2,2-Bipyridine-nickel complexes as catalysts for electrosynthesis of symmetrical ketones from organic halides (L. Garnier, Y. Rollin, J. Périchon), (367) 347

Chalcogens, contrasting roles in synthesis of mixed metal complexes (CO)$_6$Fe$_2$(μ_3-E)$_2$M(PPh$_3$)$_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

Nickel complexes containing phosphorus(III) ligands and their related Ziegler catalysts, electrochemical studies of (A.A. Pozdeeva, U.M. Dzhemilev, N.R. Popod'ko, R.I. Khusnutdinov, S.I. Zhdanov, G.A. Tolstikov), (367) 205

Oligomerisation of ethene with catalysts exhibiting enzyme-like activities (S.J. Brown, A.F. Masters), (367) 371

Organometallic complexes, extent and direction of optical induction (T. Bartik, I. Gerdes, P. Heimbach, H.-G. Schulte), (367) 359

(Tmeda)Ni(H$_2$C=CHCOOCH$_3$)$_2$, synthesis and structure of (W. Kaschube, K.R. Pörschke, K. Seevogel, C. Krüger), (367) 233

Osmium

Aromatic olefinic ruthenium(0) and osmium(0) complexes, C$_6$Me$_6$Ru(CO)(C$_2$H$_3$R) and C$_6$H$_6$Os(L)(C$_2$H$_4$), direct reductive synthesis of (H. Werner, K. Roder), (367) 339

Hydrido(methyl)osmium compounds, C$_6$H$_6$OsH(CH$_3$)(PR$_3$), synthesis, properties and reactivity of (K. Roder, H. Werner), (367) 321

Palladium

Chalcogens, contrasting roles in synthesis of mixed metal complexes (CO)$_6$Fe$_2$(μ_3-E)$_2$M(PPh$_3$)$_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

Methyl 4-tri-n-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Montmorillonitesilylaminepalladium(II) chloride, selective hydrodebromination of aryl bromides by (Y.V. Subba Rao, K. Mukkanti, B.M. Choudary), (367) C29

Oxidative addition–reductive elimination reactions; crystal structure of Me$_2$(*p*-BrC$_6$H$_4$CH$_2$)Pd(phen)Br (A.J. Canty, A.A. Watson, B.W. Skelton, A.H. White), (367) C25

Palladium(0)-mediated reactions of BrZnCH(CH$_3$)COO^{t}Bu with vinyl and aryl triflates, influence of catalyst on (F. Orsini, F. Pelizzoni, L.M. Vallarino), (367) 375

Phosphorus

Aromatic olefinic ruthenium(0) and osmium(0) complexes, C$_6$Me$_6$Ru(CO)(C$_2$H$_3$R) and C$_6$H$_6$Os(L)(C$_2$H$_4$), direct reductive synthesis of (H. Werner, K. Roder), (367) 339

t-Butyl-substituted tin–phosphor compounds, preparation and structure of (D. Hänssgen, H. Aldenhoven, M. Nieger), (367) 47

Hexahydrido(η^5-pentamethylcyclopentadienyl)rhenium(VII), (η^5-C$_5$Me$_5$)ReH$_6$, reactivity of (W.A. Herrmann, H.G. Theiler, E. Herdtweck, P. Kiprof), (367) 291

Hydrido(methyl)osmium compounds, C$_6$H$_6$OsH(CH$_3$)(PR$_3$), synthesis, properties and reactivity of (K. Roder, H. Werner), (367) 321

β-Keto phosphines derived from ferrocene; syntheses and structures of [Ph$_2$PCH$_2$C(O)(η^5-C$_5$H$_4$)Fe(η^5-C$_5$H$_5$)](L^1) and *trans*-[PdCl$_2$L$^1_2$] (P. Braunstein, T.M. Gomes Carneiro, D. Matt, F. Balegroune, D. Grandjean), (367) 117

Ligand-centred coupling of alkynyl radical cations $[Mo(C{\equiv}CR)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^+$ (R = Ph or Bu^n): crystal structure of $[Mo_2(Ph_2PCH_2CH_2PPh_2)_2(\eta\text{-}C_7H_7)_2(\mu\text{-}C_4R_2)][PF_6]_2$ (R = Ph) (R.L. Beddoes, C. Bitcon, A. Ricalton, M.W. Whiteley), (367) C21

Molybdenum(II) and tungsten(II) cationic seven-coordinate complexes, synthesis of (P.K. Baker, M. Bamber, G.W. Rogers), (367) 101

Nickel complexes containing phosphorus(III) ligands and their related Ziegler catalysts, electrochemical studies of (A.A. Pozdeeva, U.M. Dzhemilev, N.R. Popod'ko, R.I. Khusnutdinov, S.I. Zhdanov, G.A. Tolstikov), (367) 205

Organometallic complexes, extent and direction of optical induction (T. Bartik, I. Gerdes, P. Heimbach, H.-G. Schulte), (367) 359

Palladium complexes as catalyst in reduction of aromatic acyl fluorides (R. Braden, T. Himmler), (367) C12

Ruthenium dinuclear cluster with N- and O-containing bridges, reactions with CO_2 of (F. Neumann, G. Süss-Fink), (367) 175

Thiolate-amine complexes $[Ru(CO)_2(bmae)]$ and $[Ru(CO)_2(^{bu}bmae)]$ ($bmae^{2-}$ = 1,2-bis(2-mercaptoanilino)ethane(−2); ^{bu}bmae = 1,2-bis(2-mercapto-3,5-di-t-butylalinino)ethane(−2)), UV photolysis of (D. Sellmann, O. Käppler, F. Knoch), (367) 161

Trimethylphosphinecobalt(I) complexes: reaction with cyclopentatriene and cyclooctatetraene; crystal structure of $[Co(\eta^4\text{-}C_7H_8)(PMe_3)_3]BPh_4$ (L.C. Ananias de Carvalho, M. Dartiguenave, Y. Dartiguenave), (367) 187

Platinum

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(CO)_6Fe_2(\mu_3\text{-}E)_2M(PPh_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

Rhenium

Four isomeric (α-diimine)$Re(CO)_3Hal$ complexes, synthesis, electrochemistry and emission spectroscopy in solution of (W. Kaim, H.E.A. Kramer, C. Vogler, J. Rieker), (367) 107

Hexahydrido(η^5-pentamethylcyclopentadienyl)rhenium(VII), $(\eta^5\text{-}C_5Me_5)ReH_6$, reactivity of (W.A. Herrmann, H.G. Theiler, E. Herdtweck, P. Kiprof), (367) 291

Rhodium

$(\eta^5\text{-}C_5Me_5)_2CoRh(\mu\text{-}CO)_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197

Dicyclopentadienyltricarbonyldirhodium, search for ^{13}C equilibrium isotope effects in (S. Berger, M. Ochs), (367) 343

Ruthenium

Aromatic olefinic ruthenium(0) and osmium(0) complexes, $C_6Me_6Ru(CO)(C_2H_3R)$ and $C_6H_6Os(L)(C_2H_4)$, direct reductive synthesis of (H. Werner, K. Roder), (367) 339

Ruthenium dinuclear cluster with N- and O-containing bridges, reactions with CO_2 of (F. Neumann, G. Süss-Fink), (367) 175

Thiolate-amine complexes $[Ru(CO)_2(bmae)]$ and $[Ru(CO)_2(^{bu}bmae)]$ ($bmae^{2-}$ = 1,2-bis(2-mercaptoanilino)ethane(−2); ^{bu}bmae = 1,2-bis(2-mercapto-3,5-di-t-butylalinino)ethane(−2)), UV photolysis of (D. Sellmann, O. Käppler, F. Knoch), (367) 161

Selenium

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(CO)_6Fe_2(\mu_3\text{-}E)_2M(PPh_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

$(\eta^5\text{-}C_5Me_5)_2CoRh(\mu\text{-}CO)_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197

Group 6 metal tetracarbonyl complexes, *cis*-$[M(CO)_4\{(C_5H_4ECH_3)_2Fe\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1′-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Silicon

$Et_3SiCF{=}CFM$ (M = Li, ZnBr), preparation and reactivity of (P. Martinet, R. Sauvêtre, J.-F. Normant), (367) 1

Palladium complexes as catalyst in reduction of aromatic acyl fluorides (R. Braden, T. Himmler), (367) C12

Silanol silylation by a chlorosilane in the presence of neutral nucleophiles, mechanism of (S. Rubinsztajn, M. Cypryk, J. Chojnowski), (367) 27

Tetrakis- and hexakis-(trimethylsilyl)cobaltocenium ions, ring rotation hindrance in (J. Okuda), (367) C1

Sulphur

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(CO)_6Fe_2(\mu_3\text{-}E)_2M(PPh_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

$(\eta\text{-}C_5H_5)_2Ti(CO)_2$, reaction with sulphoxides of (T.L. Chen, A. Shaver, T.H. Chan), (367) C5

$(\eta^5\text{-}C_5Me_5)_2CoRh(\mu\text{-}CO)_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197

Dicarbonylcyclopentadienylmethylthiolatoiron(II), reactions with monocyanoacetylene of; crystal structures of $[FeC(O)C(CN){=}CH(SCH_3)(\eta^5\text{-}C_5H_5)(CO)]$ and of $[\mu\text{-}\eta^2\text{-}Fe(C{=}CCN)(\eta^5\text{-}C_5H_5)(CO)_2][Co(CO)_3]_2$ (R. Kergoat, M.M. Kubicki, L.C. Gomes de Lima, H. Scordia, J.E. Guerchais, P. L'Haridon), (367) 143

Group 6 metal tetracarbonyl complexes, *cis*-$[M(CO)_4\{(C_5H_4ECH_3)_2Fe\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1′-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Thiolate-amine complexes $[Ru(CO)_2(bmae)]$ and $[Ru(CO)_2(^{bu}bmae)]$ ($bmae^{2-}$ = 1,2-bis(2-mercaptoanilino)ethane(−2); ^{bu}bmae = 1,2-bis(2-mercapto-3,5-di-t-butylalinino)ethane(−2)), UV photolysis of (D. Sellmann, O. Käppler, F. Knoch), (367) 161

Tellurium

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(CO)_6Fe_2(\mu_3\text{-}E)_2M(PPh_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

Tin

Bis[8-(dimethylamino)1-naphthyl-*C,N*]tin(II) and its $W(CO)_5$ adduct, synthesis and characterization of (J.T.B.H. Jastrzebski, P.A. van der Schaaf, J. Boersma, G. van Koten, D. Heijdenrijk, K. Goubitz, D.J.A. de Ridder), (367) 55

t-Butyl-substituted tin–phosphor compounds, preparation and structure of (D. Hänssgen, H. Aldenhoven, M. Nieger), (367) 47

Methyl 4-tri-n-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Titanium

Bis(cyclopentylcyclopentadienyl)titanium dichloride and bis(cyclohexylcyclopentadienyl)titanium dichloride, crystal structures of (M. Kapon, G.M. Reisner), (367) 77

$(\eta\text{-}C_5H_5)_2Ti(CO)_2$, reaction with sulphoxides of (T.L. Chen, A. Shaver, T.H. Chan), (367) C5

Dicyclopentadienyltitanium dihalides, electronic structure as indicated by UV photoelectron spectrocopy of (T. Vondrák, K. Mach, V. Varga), (367) 69

Tungsten

Bis[8-(dimethylamino)1-naphthyl-*C,N*]tin(II) and its $W(CO)_5$ adduct, synthesis and characterization of (J.T.B.H. Jastrzebski, P.A. van der Schaaf, J. Boersma, G. van Koten, D. Heijdenrijk, K. Goubitz, D.J.A. de Ridder), (367) 55

Group 6 metal tetracarbonyl complexes, *cis*-$[M(CO)_4\{(C_5H_4ECH_3)_2Fe\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1′-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Molybdenum(II) and tungsten(II) cationic seven-coordinate complexes, synthesis of (P.K. Baker, M. Bamber, G.W. Rogers), (367) 101

Vanadium

Cyclopentadienylalkynevanadium(I) CO-free complexes, facile synthesis and reactivity of (B. Hessen, J.H. Teuben), (367) C18

Zinc

$Et_3SiCF{=}CFM$ (M = Li, ZnBr), preparation and reactivity of (P. Martinet, R. Sauvêtre, J.-F. Normant), (367) 1

Palladium(0)-mediated reactions of $BrZnCH(CH_3)COO^tBu$ with vinyl and aryl triflates, influence of catalyst on (F. Orsini, F. Pelizzoni, L.M. Vallarino), (367) 375

Zirconium

Boron hydride insertion into Zr–C bond of dimeric (η^2-formaldehyde)zirconocene (G. Erker, U. Hoffmann, R. Zwettler, G. Krüger), (367) C15

$(C_5H_5)_2ZrCl(AlCl_4)$, crystal and molecular structure and unusual hydrocarbon solubility properties of; relevance to Ziegler–Natta polymerization chemistry (M.V. Gaudet, M.J. Zaworotko, T.S. Cameron, A. Linden), (367) 267